



POSTER PRESENTATION

Open Access

An alternative analgesic agent for intra-articular corticosteroid injections (IAI) in juvenile idiopathic arthritis (JIA) – a fentanyl lollipop

Z Zareen*, D Finn, O Killeen

From 18th Pediatric Rheumatology European Society (PReS) Congress
Bruges, Belgium. 14-18 September 2011

Background

Intra articular injections are an effective and widely used technique that involves the direct application of a corticosteroid compound such as triamcinalone hexacetonide into a joint with active synovitis. Local IAI's can avoid many of the unwanted adverse effects of systemic steroids but in children younger than 11 years or those with a polyarticular presentation a general anaesthetic is typically required. Traditional local anaesthetic agents include topical agents such as lignocaine or gaseous agents such as entonox (nitrous oxide). We have introduced an unique form of analgesia for this procedure since the establishment of our national unit in 2006 in form of fentanyl lollipops (Actiq).

Aim

To evaluate the efficacy & safety of fentanyl lollipops for children with JIA undergoing IAI's.

Methods

A total of 19 patients were studied. Pain scores an hour after the procedure were recorded using a 0–10 cm visual analogue scale (VAS). Parents and patients were reinterviewed at a later date regarding the effectiveness of the lollipops as a form of pain relief and whether they would consider this form of analgesia for their child again.

Results

A total of 25 joints were injected in 19 patients (median age 14.6 yrs, range 10.4–16.6 yrs). The median patient and

parent pain scores were 1.5 and 4.0 respectively. There were no serious adverse events in any patient. All parents agreed that fentanyl lollipops were safe and effective, and over 90% would opt for this form of analgesia again.

Conclusion

Fentanyl lollipops are a safe and an effective form of analgesia for intra-articular injections in children. A significant number of parents overestimated pain related to the procedure compared to their child.

Published: 14 September 2011

doi:10.1186/1546-0096-9-S1-P205

Cite this article as: Zareen et al.: An alternative analgesic agent for intra-articular corticosteroid injections (IAI) in juvenile idiopathic arthritis (JIA) – a fentanyl lollipop. *Pediatric Rheumatology* 2011 **9**(Suppl 1):P205.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: zuneraz@yahoo.ie

National Centre for Paediatric Rheumatology, Our Lays Children Hospital, Crumlin, Dublin, Ireland